- **James Byrne, M.D., Ph.D.** – 2023 Awardee

  Assistant Professor
  University of Iowa

  “Novel materials to improve management of hypoxic tumors ”

Malignant peripheral nerve sheath tumors (MPNSTs) represent a devastating type of tumor that accounts for 5 –10% of all soft-tissue sarcomas and has a 5-year overall survival rate of only 50%. Despite curative-intent treatment (surgery and/or radiotherapy) of localized MPNSTs, prognosis remains poor. The objective of the proposed research is to overcome these barriers by taking advantage of a novel series of GeMs that we recently developed to safely and tunably release gases. GeM manufacturing utilizes a simple, cost-effective and scalable technology involving components FDA classified as Generally Regarded As Safe (GRAS), and we propose to develop GeMs to safely and sustainably deliver oxygen through intratumoral administration. We aim to demonstrate that mitigating tumor-associated hypoxia using intratumorally administered O2-GeMs will improve responses to radiation and immunotherapy in preclinical models of MPNSTs and is safe for use in MPNST patients undergoing neoadjuvant radiation therapy.

- **Milan Chheda , M.D.** – 2023 Awardee

  Associate Professor of Medicine and Neurology

- **Agnieszka Czechowicz , M.D., Ph.D.** – 2023 Awardee

  Assistant Professor of Pediatrics
  Stanford University

  "Development of Novel DNA - guided Argonaute Base Editing"

Approximately half of the known human genetic diseases are caused by single- nucleotide variants (SNVs). While CRISPR-base editing (BE) technologies offer hope in the treatment of SNV-related diseases, they can only edit certain regions of the genome and lack perfect specificity. To overcome these limitations and enable exact correction of all SNVs, we propose to develop a novel DNA - guided Argonaute (Ago) BE technology by using Ago - deaminase fu sion protein as Ago base editor, facilitated by the sequence- specific strand invasion function of the peptide nucleic acid (PNA).

We will first demonstrate targeted deamination by Ago base editor (Aim 1), and then successful BE combining Ago base editor and PNA (Aim 2). The optimized Ago BE system will then be applied to correct pathogenic mutations causing severe combined immunodefi ciency (SCID) in fibroblasts (Aim 3), and subsequently in mouse hematopoietic stem cells (HSCs) (Aim 4). These edited HSCs will then be transplanted to cure disease in a SCID mouse model. The feature of limitation - free binding of Ago and PNA renders this technology unprecedented applicability, and this dual binding mechanism further guarantees its accuracy. A highly versatile, accurate BE system could lead to curative treatments for millions of individuals with various disease - causing SNVs.

- **Pankaj Desai, Ph.D.** – 2023 Awardee

  Professor and Chair, Division of Pharmaceutical Sciences
  University of Cincinnati

  "Clinical and Correlative Translational Studies for Repurposing Letrozole as a Novel Therapeutic for Glioblastoma: Proof of Efficacy in Combination with Metronomic Temozolomide "

Our goal is to expedite repurposing letrozole, (LTZ), an aromatase inhibitor widely used for breast cancer treatment, as a novel drug for glioblastoma (GBM), a lethal brain tumor. The objective of this proposal is to conduct a "proof-of-efficacy" study as a critical first-step to support our commercialization efforts and facilitate larger confirmatory phase II trials. Our observations include: 1) the enzyme aromatase is abundantly expressed in GBM, 2) LTZ readily crosses the blood-brain barrier and exerts marked anti-GBM activity in pre-clinical models, 3) LTZ synergistically potentiates the activity of temozolomide (TMZ), the only drug approved for GBM, and, 4) in a phase 0/1 study in recurrent GBM patients (dose range, 2.5-15 mg), the 15 mg dose was identified as the Recommended Phase 2 Dose (RP2D) since it met the primary endpoint (tumoral concentration > 2 μM) with no safety concerns. Based on the Simon two-stage optimal study design, here we propose to initiate stage one efficacy study in recurrent GBM patients (N = 19) who will receive LTZ (15 mg) with metronomic TMZ (50 mg/m2). A 30% improvement in 6-month progression-free survival will meet the criteria to reject futility, confirm efficacy and establish RP2D of LTZ for further development.

- **Bryan Dickinson , Ph.D.** – 2023 Awardee

  Assistant Professor of Chemistry
  University of Chicago

  “Development of targeted translational activating oligos to treat SYNGAP1 deficiency - related neurological disorders ”

- **Andrea Domenighetti , Ph.D.** – 2023 Awardee

  Research Scientist and Principal Investigator
  Rehabilitation Institute of Chicago

  “Repurposing 5-azacytidine for the treatment of muscle contractures in children with cerebral palsy.”

- **Amir Hajrasouliha, M.D.** – 2023 Awardee

  Assistant Professor of Ophthalmology
  Indiana University

  "A Novel Minimally - Invasive Vision Restoration Method Through Continuous Slow - Release of Artificial Photoreceptors "

We have devised an innovative solution for vision loss affecting 250 million people worldwide. The root cause of the problem is deterioration of retinal photoreceptors in retinal dystrophies and degenerations. The diversity and complexity of different gene s involved in these diseases have made photoreceptor replacement therapy a complex clinical challenge. We have identified a potential candidate in an artificial photoreceptor termed nanocomposite, capable of restoring sensory functions on the retina of bli nd mice model (rd1) and in ex vivo retina explants. However, these nanocomposites have a limited lifespan and must be renewed and replaced for the eye to maintain visual resolution. In this project, we aim to devise a method to continuously release our pro

- **David Ingbar , M.D.** – 2023 Awardee

- **James Mathew , Ph.D., M.Sc.** – 2023 Aww 00bc0eBnee

- **Hiromitsu Nakauchi, M.D., Ph.D.** – 2023 Awardee

  Professor of Genetics
  Stanford University

  “Metabolic Pausing: Revolutionizing Affordable Organ Transplantation Globally ”

Addressing the global organ shortage, our long- term goal is to economically generate universally immune - compatible organs that are readily available for transplantation, democratizing access to this life - saving resource and procedure. Liver diseases alone cause

i12.9 tyh41.7 ( )0.6 lor ai32 (e)]TJ -0.002 Tc -0.007 Twnt ating for orga73.2 (n)1.8 ( )0.5 (t)-6 (r)-7.9 (a)7.2 (n)1.7 (s)-5.2 (p)
i 6 1 ( ) ( ) 0 6 1 ( ) ( ) 1 8 ( ) 8.7 . 1 ( ) 1 ( ) 1 ( )

- **Sean Pitroda , M.D.** – 2023 Awardee

  Associate Professor of Radiation and Cellular Oncology
  University of Chicago

  “Validation of Tumor Aneuploidy as a Novel Predictor of Radio - Immunotherapy Response in Metastatic Non - small Cell Lung Cancer: Biomarker Analysis of a Prospective Randomized Phase I/II Clinical Trial”

Although immunotherapy has revolutionized the treatment of metastatic NSCLC, only a small fraction of patients experiences long - term survival following treatment. In a recent breakthrough, we discovered that a subset of patients with immunotherapy resistan t, highly aneuploid NSCLC derive a major benefit from the addition of radiotherapy to immunotherapy leading to improved outcomes. Here, we propose to c -0.016 Tw 0.5 0 Td [a pa nov al198 (s9 (o)19.5 (

- **Carlos Subauste, M.D.** – 2023 Awardee

  Professor of Medicine and Pathology
  Case Western Reserve University

  "CD40-TRAF2,3 Blocking Peptide for the Treatment of Ocular Disorders"

The main cause of blindness in patients with diabetic retinopathy is macular edema caused by vascular leakage. Anti-VEGF agents are used to treat diabetic macular edema. Unfortunately, many patients do not respond because vascular leakage is also driven by VEGF-independent inflammation. The objective of this application is to identify an improved approach to treat diabetic retinopathy. We hypothesize that, in contrast to anti-VEGF agents, nanoparticle-driven sustained intravitreal release of a peptide that blocks CD40-TRAF2,3 signaling will cause long-lasting reduction of VEGF and inflammatory molecules, reversing VEGF-dependent and independent vascular abnormalities (Claudin 5 redistribution) and reducing vascular leakage in diabetic mice. Using confocal microscopy and mass spectrometry, we will first establish how long the peptide is present in the retina and inhibits CD40. In the second aim, we will establish nanoparticle-assisted method of sustained delivery of the peptide and determine its duration of action in the retina. Using a mouse model of diabetic retinopathy in the third aim, we will test whether nanoparticle-associated blocking peptide is more effective than an anti-VEGF agent in reducing expression of inflammatory molecules, VEGF, vascular leakage and normalizing Claudin 5 expression. This work may result in a novel and improved approach to treat diabetic retinopathy.

- Paul Turner , Ph.D.

- **Richard Vile, Ph.D.** – 2023 Awardee

  Professor of Immunology
  Mayo Clinic

  "In Vivo Generation of Persistent, Memory and Effector CAR T Cells from Naïve T Cells as a Platform for Treating Solid Tumors "

Activation of naïve T cells by pathogens induces differentiation into memory T cells that can both produce and replenish the supply of effector cytotoxic T cells. CAR T cells (CAR T) are engineered to express a Chimeric Antigen Receptor (CAR) which directs the T cell to kill tumor cells. Clinically produced CAR T primarily contain effector cells for tumor killin g with few replenishing memory CAR T, leading to tumor escape once effector CAR T die. We engineered dendritic cells (DC) ex vivo to release a CAR vector during presentation of an immunogenic epitope to naïve T cells. Activation of naïve T cells is coupled with infection of T cells by the CAR vector, producing populations that differentiate into effector CAR T (to kill tumor) and memory CAR T to replace lost effectors. We hypothesize that direct in vivo administration of a DC - targeting vector will enable DC - based generation of both CAR T populations in situ from naïve T cells. We will (1) compare vectors for delivery of CAR/packaging functions to DC, T cell targeted CAR vector release in vivo, and potential toxicities; (2) test anti - tumor efficacy in vivo and (3) identify IND - enabling studies to prepare for IND application to clinical trial.

Robert Abramovitch, Ph.D. – 2022 Awardee

Associate Professor of Microbiology and Molecular Genetics
Michigan State University

"Development of New MmpL3 Inhibitors to Treat Mycobacterial Infections"

Tuberculosis (TB) remains a global health crisis causing ~1.5 million deaths per year. To cure TB, treatment requires daily, multidrug therapy for 6 months. However, with the evolution of drug resistant Mycobacterium tuberculosis (Mtb), current therapies a re inadequate to control multi- drug resistant TB.

MmpL3 inhibitors are attractive new agents functioning against drug susceptible and resistant Mtb. We have discovered a new class of MmpL3 inhibitors called HC2099. We optimized and found an orally bioavailable analog that is efficacious in vivo in an acut e murine Mtb infection model. However, we encountered a barrier to development caused by in vivo metabolism of HC2099, leading to a short half - life. We have identified the metabolic liability presenting a clear path forward to overcome this barrier to deve lopment.

The goal of this proposal is to further optimize the HC2099 series to generate proof- of- concept in vivo efficacy data showing its utility for once daily dosing.
To fulfill these goals, we will complete:

Specific Aim 1. Define structure activity relationships and optimize activity of HC2099 series.
Specific Aim 2. Conduct integrated SAR and pharmacokinetic studies to prioritize compounds for in vivo efficacy studies.
Specific Aim 3. Demonstrate proof - of- concept in vivo efficacy of HC2099 series against Mtb infection.

Drew Adams, Ph.D. – 2022 Awardee

Associate Professor, Dept. of Genetics; Thomas F. Peterson Jr Professor of Novel Therapeutics
Case Western Reserve University

“Targeting Exportin -1 to Block T Cell Activation in Autoimmune Diseases”

My laboratory seeks to validate new targets for drug discovery, and our past work has led to high-impact publications and commercialization. Recently we identified Exportin -1 (XPO1)
1 is10.8 ( )0.2 (a)16.2 ( )0.5 (v)6.2 (a)16.9 (l)8.9 (i)8.d (F)17.2 (a)16.2 (t)ed (F)17. (e)0.d (F)17.ruF taF
-1 a[illegible]

Nita Ahuja, M.D. – 2022 Awardee

Ethan Anderson, Ph.D. – 2022 Awardee

Associate Professor of Experimental Therapeutics
University of Iowa

“Development and validation of a piezoelectric immunosensor to rapidly detect a novel biomarker of morbidity and mortality risk in sepsis patients.”

Amanda Garner, Ph.D. – 2022 Awardee

Assistant Professor of Medicinal Chemistry
University of Michigan

"Decoding the Druggable Transcriptome"

Following completion of the Human Genome Project, it was revealed that only ~2% of our genome encodes for proteins, and the overwhelming majority of our transcribed genome is comprised of often highly conserved non- coding RNAs. Through expanded exploration of RNA biology, a diversity of structure /eer (t)-16(o)-22.2 (l) 24ah (c)-1(Tc 0.00 10.08 -0 0 10.08 51s)-14.2 (i)-9.1 (n( )0

Justin Lathia, Ph.D. – 2022 Awardee

Associate Professor of Cellular and Molecular Medicine
Cleveland Clinic Lerner Research Institute

“Development of an anti-cancer stem cell therapy for glioblastoma by targeting the epigenetic state via WDR5”

Our long-term objective is to synthesize a brain penetrant, WDR5 inhibitor with sufficient potency to decrease cancer stem cell (CSC) proliferation, self-renewal, and viability. We

Craig Levin, Ph.D. – 2022 Awardee

Professor of Radiology, Physics, Electrical Engineering, and Bioengineering
Stanford University

"Multiplexed Positron Emission Tomography Imaging of Promising Biomarkers for Immune Checkpoint Inhibitor Therapy"

Immunotherapy has revolutionized medical oncology. The use of immune checkpoint inhibitors (ICIs) has demonstrated revolutionary results in cancer treatment, but only a portion of patients can achieve complete disease remission. Current diagnostic imaging based on the tumor size measurement alone is not adequate to assess initial response to immunotherapy and disease evolution. Therefore, there is an urgent need for the development of new non - invasive imaging assays that can tell whether a patient's cancer

Feng Lin, B.Sc., Ph.D.– 2022 Awardee

Professor of Molecular Medicine
Cleveland Clinic Lerner Research Institute

"Development of a New Drug for Patients With T Cell Lymphoma"

T-cell lymphomas (TCL) are complex clusters of aggressive blood cancers with unsatisfactory management options. In pilot studies, we identified CD6 as a novel therapeutic target for TCL. We developed an antibody-drug conjugate (ADC) specific for CD6 (CD6-ADC) using a conventional anti-CD6 monoclonal antibody (mAb) and found that this ADC is highly effective in selectively killing TCL cells both in vitro and in vivo. Nanobodies, heavy-chain only antibodies produced in camelids, are the next generation of mAb-based therapeutics. Given the many advantages that nanobodies have over the conventional mAbs, especially their excellent tumor penetrating capacity, we are developing the next generation of CD6-ADC using nanobody technology. We have immunized an alpaca and identified more than 120 anti-CD6 nanobody candidates. In this project, we will rigorously characterize these candidates to identify the best anti-CD6 nanobodies, use them to develop the next generation of CD6-ADC, and evaluate their treatment efficacy and potential adverse effects both in vitro and in vivo in a preclinical model of TCL. These high-risk studies, if successful, will generate the next generation of CD6-ADC and provide the required proof of concept to translate it into a much-needed new dru g for patients with TCL.

Jonathan Marchant, Ph.D., M.A. – 2022 Awardee

Professor
Medical College of Wisconsin

"IDENTIFYING NEW DRUGS TO TREAT AN INFECTIOUS DISEASE OF POVERTY"

Diseases caused by parasitic flatworms impact millions of people. These infections cruelly target some of the most disadvantaged, with an especial burden on children worldwide. Many infections are treated with a drug called praziquantel (PZQ), the sole cli nical agent available, but not an ideal treatment: for 40 years we have not understood how PZQ works, and it is not effective against every parasitic disease. Therefore, discovery of new anti - parasitics is a priority.

Our team recently identified the target of PZQ, overcoming a long - standing roadblock. The target is an ion channel, named TRPMPZQ, responsible for PZQ - dependent worm paralysis and elimination. Our discovery explains why PZQ is effective against some diseas es and not others, as the TRPMPZQ binding site differs between parasites. With this knowledge in hand, we are optimally placed to discover novel drugs.

Therefore, in this 'Catalyst' proposal, we will take two independent, but parallel approaches to identifying novel chemotypes active at TRPMPZQ. We will take a rational drug engineering approach (Aim 1) and an unbiased drug screen (Aim 2). By adhering to 1 0 quantitative metrics that benchmark success of our assays, we aim within 18 months to discover novel agents that can be further progressed in subsequent work.

Tay Netoff, Ph.D. – 2022 Awardee

Professor of Biomedical Engineering
University of Minnesota

Arjun Raman, M.D., Ph.D. – 2022 Awardee

Assistant Professor
University of Chicago

"The 'statistical' design of synthetic microbiomes"

Complex microbial communities ('microbiomes') are ecosystems that critically contribute to human physiology. A major goal is to design such communities to unlock their immense potential in addressing challenges affecting human health. However, it is remark ably difficult to rationally engineer microbiomes due to their immense complexity —the number of possible interactions between the thousands of component microbes is astronomical. Thus, what strategy can be used to design functional microbiomes? We have dev eloped a 'top - dowi-5 ( )0f16.ti(t)3 (iy1(es)3..1 (y)]TJ 0 1.048 0 Ta)-4.2 5 Tw T*tmroyi

p

Jeffrey Schneider, Ph.D. – 2022 Awardee

Assistant Professor
Rush University Medical Center

"Glycoengineering Anti-HIV antibodies to increase delivery to the brain and block viral egress from the CNS"

There is no cure for HIV despite great strides in combination anti-retroviral therapy(cART) to control HIV/AIDS. Once cART is interrupted, HIV re-emerges from sanctuary sites to reseed the body. One such site is the central nervous system (CNS). Our recent chimeric human brain mouse model demonstrates that HIV egresses from the brain to peripheral organs following cART cessation. Therefore, it is critical to develop therapies targeted to the brain to block this egress if a viable cure strategy is going to be achieved. We propose to target anti-HIV neutralizing antibodies to the brain to block this egress. CNS antibodies have a distinct glycosylation profile to support their retention in the CNS in comparison to bulk IgG. We propose a novel methodology to target antibodies to the brain, involving engineering antibodies to be more "CNS like" through glycosylation and subclass manipulation (Aim 1). We will assess these "CNS like" anti-HIV antibodies for CNS penetration and potential to block HIV viral egress following cART cessation in our chimeric human brain mouse model (Aim 2). Collectively, these studies will explore the validity of engineering CNS-like antibodies to increase penetration to the brain and ability to block viral egress following cART cessation.

Jennifer Woyach, M.D. – 2022 Awardee

Assistant Professor of Internal Medicine
The Ohio State University Wexner Medical Center

"Optimizing BTK Inhibitor Therapy in CLL"

Chronic lymphocytic leukemia (CLL) is the most prevalent adult leukemia. Covalent Bruton's Tyrosine Kinase inhibitors (BTKi) ibrutinib and acalabrutinib have been paradigm - shifting in

Xiaoyu Zhang, Ph.D. – 2022 Awardee

Assistant Professor of Chemistry
Northwestern University

"Discovering and developing neoantigen inducers as new and effective cancer immunotherapeutics"

Cancer immunotherapy relies on cytotoxic T lymphocytes (CTLs) to recognize proteins or mutation - derived peptides displayed on cancer cell surfaces in order to eliminate cancer cells. Mutation - derived peptides, or neoantigens, that display on major histocom patibility complex I (MHC - I) on cancer cell surfaces are particularly relevant to effective cancer immunotherapy. Although exome sequencing and in silico algorithms are used to predict MHC - I-

Swetha Gowrishankar , Ph.D. - 2021 Awardee

Anatomy and Cell Biology
University of Illinois College of Medicine at Chicago

"Evaluation of the In Vivo Efficacy of Novel Autophagy Activators for Amelioration of Alzheimer's Disease Pathology in the 5xFAD Mouse Model "

Alzheimer's disease (AD) is a progressive neurodegenerative disease that affects over 5 million people in the United States and is characterized by amyloid plaques, tau tangles, loss of synapses and ultimately, neuronal death. FDA - approved drugs for AD tre at the symptoms of the disease but do not improve the underlying cell damage that leads to disease progression, highlighting the need for novel neuroprotective therapeutic options. Dysfunction in the autophagic and lysosomal pathways, which are critical fo r protein and organelle homeostasis in neurons, is associated with different stages of the disease, and thus modulation of autophagy has emerged as a new strategy for developing AD - targeted drugs. To this end, we have identified novel, small - molecule autop hagy activators in a high - throughput screen that rescued key pathological features in human neuronal culture models of the disease. F-8 (e)( )0.6 (f)3 (i)-7 (v)-9.r.( )0.6 (.)0.5 (F-8 0 Tw 3) (u)13.3.1 (h)68 (e)( )614.2 ( )0.54 Td [(002 Tw 0.

Jonathan Kurtis , M.D., Ph.D. - 2021 Awardee

Pathology and Laboratory Medicine
Brown University

"Apoptosis - inducing Anti - Malaria Drugs Targeting PFGARP "

The overall aim of this application is to discover novel therapeutics for Plasmodium falciparum malaria. P. falciparum is a leading cause of morbidity and mortality in developing countries, infecting hundreds of millions of individuals and killing over 300 ,000 children each year 1. The spread of parasites resistant to the artemisinin family of compounds 2 threatens recent progress achieved by antimalarial campaigns and underscores the urgent need to identify new anti - malarial drugs.

In previous work, we discovered PfGARP, a previously unrecognized vaccine candidate found only in P. falciparum. Antibodies to the highly invariant carboxyl- terminal of PfGARP (PfGARP - A, aa 411673) inhibit parasite growth in vitro by 99% compared to contr ols ($P < 0.001$) by killing trophozoite stage parasites. In confocal and transmission electron microscopy studies, PfGARP localized to the exofacial surface of the RBC membrane in trophozoite and early schizont infected RBCs, but not to other parasite stages or uninfected RBCs. Importantly, the growth inhibition assays are performed in the absence of any immune effector molecules (complement) or cells - thus the remarkable anti - parasite effect of anti - PfGARP results from antibody binding alone. This is furthe r supported by the killing effect of recombinant mAb (KD 2.9 nM, (95% CI = 1.3 – 5.9 nM)) and its rec monovalent Fab that target aa 443 - 459 (VKNVIEDEDKDGVEIIN) of PfGARP.

The Scientific Premise of this application is that PfGARP is a high - value druggable target based on: 1) its surface expression on infected RBCs, 2) the absence of any significant amino

Kelvin Lee, M.D. - 2021 Awardee

Medicine
Indiana University

"Targeting PIM2 and its Regulation of the c - Myc Oncogene in Multiple Myeloma and Other Cancers"

Multiple myeloma (MM) is a malignancy of plasma cells that is the second most common hematologic malignancy (20% of all cases), and remains incurable for almost all patients. The primary cause of treatment failure is upregulation of pro - survival resistance mechanisms, and identifying these mechanisms remains key for new therapeutic development. We have recently reported that the serine- threonine kinase PIM2 has a major pro - survival role in MM, and that inhibition of PIM2 with a first - in - class PIM2- selective non - ATP - competitive kinase inhibitor JP11646 (JP) caused significant MM cell death in vitro and in vivo in preclinical MM models. Unexpectedly, JP was much more effective than the ATP - competitive PIM kinase inhibitors due to JP's unique ability to downreg ulate of PIM2 gene and protein expression. This has led to our findings that PIM2 has critical but previously unrecognized kinase - independent (KI) functions, which would be a change in paradigm. One such PIM2 KI function that is inhibited by JP is inductio n of the expression (mRNA and protein) of the c - Myc oncogene, which is dysregulated in ~50% of all human cancers, including MM. Our additional findings suggest a completely novel mechanism where JP disrupts PIM2 interaction with a partner protein (possibly c- Myc itself) that results in downregulation of c - Myc expression and loss of c - Myc driven PIM2 gene expression, collapsing a self - reinforcing loop that sustains both PIM2 and c - Myc expression. Targeting c - Myc expression through PIM2 inhibition represents an entirely unexplored therapeutic approach. The overall goal of this proposal is to develop JP11646 for the treatment of refractory/relapsed MM, and potentially for other cancers. The Specific Aims of the proposal are:

Aim 1. Define the kinase- independent mechanisms by which PIM2 supports MM survival, and how JP11646 inhibits these.

Aim 2. Conduct preclinical and IND enabanwrnenaban i0 T65 (w)1.(x)9.4 (r)8c3 (k)-56.4 (l8 (el)8.r (6)2r2 (r-4.2 (r)s)3.9)

Tristan Maerz, Ph.D. - 2021 Awardee

Orthopaedic Surgery
University of Michigan

"Targeting Pathological Rspo2 - mediated Wnt/ β- Catenin Signaling as a Novel Treatment for Osteoarthritis "

oacti91.1 so Re

Reshmi Parameswaran, Ph.D. - 2021 Awardee

Medicine
Case Western Reserve University

"BAFF CAR-NK Cells: An Efficacious and Safe Immunotherapy for B Cell Cancers"

CAR-T immunotherapies have produced remarkable clinical responses, but several challenges remain including disease relapse due to antigen escape/decreased CAR-T persistence, fatal side effects (cytokine release syndrome, neurotoxicity), the cost and time to produce CAR-T cells. We have developed novel B cell activating factor (BAFF) ligand-based BAFF CAR with multi-receptor specificity to target BAFF receptors (BCMA, TACI, BAFF-R) expressed by B cell cancers. We deliver BAFF-CAR to human NK cells using a non-viral TcBuster transposon platform, enabling safer, simpler, and more cost-effective CAR-

Richard Pomerantz , Ph.D. - 2021 Awardee

Biochemistry
Jefferson Medical College of Thomas Jefferson University

"PROTACs for Targeting BRCA - Deficient Cancers "

DNA polymerase (Polq) is a unique polymerase- helicase DNA repair protein that has recently been validated as a synthetic lethal drug target in cancer cells harboring mutations in BRCA1 or BRCA2 (BRCA) tumor suppressor proteins. Polq promotes DNA repair via the microhomology - mediated end - joining (MMEJ) pathway (also referred to as alternative end - joining). Polq is essential for BRCA - deficient cells, but is dispensable for normal cells and Polq null mice show no phenotypes. Thus, Polq is synthetic lethal with BRCA1/2. Consistent with this, recently published Polq inhibitors (Polqi) selectively kill BRCA - deficient cells, while showing no effects in BRCA - proficient cells. Yet, whether these early stage Polqi will lead to clinically effective drugs is unknown.

Towards the g 85joso c, l.62[( )0.2f -1.214 Td [(c)7.2 (lin)1.8 (ic)7.h12.9 (h) (e)-9.42 (lly)-2.9 (r)-15 (e)fer at9.42 ((h)-5.3

-

Wei Xu, Ph.D. - 2021 Awardee

Oncology
University of Wisconsin - Madison

"CARM1 Inhibition Enhances Immunotherapy Response in Triple - Negative Breast Cancer"

Triple - negative breast cancer (TNBC) is the most aggressive breast cancer. Treatment options for this subtype are limited, as it is lacks expression of ER, PR, and HER2. Immune checkpoint inhibitors (ICI) have shown promising effects, but response rates ar e low, underscoring the need for combination therapy. However, there are many critical roadblocks for developing combination regiments, including identifying which combinations will be effective, and identifying biomarkers to select patients that are most likely to benefit from treatment.

This project is based on our groundbreaking discovery of the role of CARM1 - mcwe/16.2(n)10.8 (ee(t)3.1 (07 Tc -(t)3 0

Hongwei Yu, M.D. - 2021 Awardee

Anesthesiology
Medical College of Wisconsin

"A Novel Analgesic Approach for Chronic Pain: Small Peptide Inhibition of Nav1.7 in Anatomically Targeted Sensory Neurons "

The peripheral nervous system is an appealing site for devising new analgesic treatments

“Lupus Nephritis Biomarkers for the Advancement of Therapies for Lupus Nephritis”

Lupus, incurable autoimmune disease affects round 1.5 million persons living in the United States. Among the main risk factors for poor disease outcome is the presence of lupus nephritis (LN). The applicants have discovered and initially validated a panel of LN-biomarkers [NGAL,MCP-1,Kim-1,ceruloplasmin,adiponectin,hemopexin]. Considering the urine levels of these LN-biomarker we delineated an algorithm, the Renal Activity Lupus Index (RAIL), where higher scores reflect more kidney inflammation as seen on kidney biopsy. Changes in RAIL scores allow us to predict LN flares and LN response to therapy at least 3 months earlier than when using current clinical tools. Further, our pilot studies suggest that the RAIL score can be used to refine the dosing of current LN therapies; we observed that high-dose intravenous steroids lead to a dramatic reduction of the RAIL score that does not occur when only daily oral steroids are used for LN therapy.

Critical Scientific Roadblock to be addressed: To enable the application of these highly promising LN-biomarkers and the RAIL algorithm in clinical care and for research, following work needs to be conducted: (1) development of a high-quality multiplex assay (RAIL-MPLA) to rapidly measure all of the LN-biomarkers concurrently; (2) additional validation of the RAIL in an independent cohort of adults and children.

Approach: Working together with Ethos Research & Development and pharmaceutical companies, we will: (1) develop a RAIL-MPLA; (2) define changes of RAIL scores that reflect clinically relevant improvement of LN and reference values for absent, controlled and active LN across age-ranges; and (3) test the hypothesis that high-dose intravenous steroids result in a significantly higher reduction of the RAIL score compared to standard doses of oral steroids alone.

Expected Results/Deliverables:

- Develop a novel clinically actionable multiplex assay that accurately quantifies the biomarkers used in the RAIL within 4 hours of sample receipt.
- Increase the usability of RAIL by refining the interpretation of RAIL scores
- Immediately advance the treatment of LN by providing the scientific underpinning for the preferred use of intravenous rather than oral steroids alone.

Hannelore Heemers, Ph.D. - 2020 Awardee

Cancer Biology
The Cleveland Clinic

"Developing Citron Kinase Inhibition As A Mechanistically Novel Approach To Overcome Cancer Treatment Resistance"

Treatment resistance causes nearly all cancer deaths. Novel treatments that bypass this resistance and inhibit the biology that drives cancer progression are needed to improve patient survival. Using prostate cancer (CaP) as model, we isolated citron kinase (CIT) as a target for such a therapy. CIT is a previously unrecognized novel druggable target that is functionally distinct from current targets yet is a critical determinant of cancer cell proliferation and induces cancer growth and treatment resistance. CIT silencing inhibited growth of cell lines and xenografts representing different stages of CaP progression and diverse forms of treatment resistance but not of benign epithelial prostate cells. CIT's stimulation of cancer progression relied entirely on its kinase activity, isolating CIT's kinase activity as a completely new target. A specific CIT inhibitor has not yet been developed. By analyzing the target spectrum of kinase inhibitors and CIT kinase assays, we identified 3 multikinase inhibitors that inhibit CIT, decrease CaP cell proliferation and inhibit CIT substrate phosphorylation. Because these drugs are not selective for CIT and thus not likely to succeed as cancer therapies, we developed a lead compound that inhibits CIT at low doses, has improved CIT selectivity and promising DMPK.

Our objective is to develop our lead compound into first-in-class selective CIT inhibitor as a novel cancer treatment. We hypothesize that novel CIT inhibitors will overcome acquired resistance to conventional cancer therapies, which will be reflected in the phosphorylation status of CIT substrates. We will test 2 Specific Aims:

1. To determine the therapeutic efficacy of novel CIT inhibitors during cancer progression using multiple chemistry and crystallography approaches and kinome screens to improve CIT specificity and DMPK and verifying growth inhibition of clinical relevant treatment-resistant cancer models.
2. To determine the substrates by which CIT kinase action conveys aggressive cancer behavior using integrated state-of-the-art biotin-based proximity ligation assays, kinase substrate arrays, and mass spectrometry approaches.

Evangelos Kiskinis, Ph.D. - 2020 Awardee

Neurology
Northwestern University Feinberg School of Medicine

“Development of Spherical Nucleic Acid - Based Anti - Sense Oligos for the Therapeutic Treatment of Pediatric Epilepsy Disorders”

Heterozygous mutations in KCNQ2, which encodes a pore- forming K+ channel subunit responsible for neuronal M - current, cause neonatal epileptic encephalopathy (EE), a complex disorder presenting with severe early - onset seizures and impa8 (u)0.8 (r)1 (e02 0.05 (wo 4702 Tc 13( )016.2

Eric Morrow, M.D., Ph.D. - 2020 Awardee

Molecular Biology, Cell Biology and Biochemistry
Brown University

"Development of a Preventative Treatment for a Novel Neurometabolic Disorder in Childhood"

Intellectual disabilities are common and carry high lifetime costs to families. We have identified a new neurodevelopmental disorder caused by loss - of - function mutations in the mitochondrial enzyme glutamate pyruvate transaminase 2 (GPT2). GPT2 disease inv olves postnatal microcephaly, cognitive disability, epilepsy, and progressive spastic paraplegia. GPT2 localizes to mitochondria and catalyzes the reversible addition of an amino group from glutamate to pyruvate, yielding alanine and alpha- ketoglutarate, a metabolite in the tricarboxylic acid (TCA) cycle. Metabolic diseases in children, such as GPT2 disease, may be amenable to treatments, via dietary restrictions or supplements, that can prevent intellectual disabilities when treated early. With the prospect of newborn screening supported by genome- wide sequencing on the horizon, there will be new opportunities to intervene in childhood brain disease. Therefore, we rapidly need to determine which new neurometabolic diseases may be amenable to interventions in early childhood. Our preliminary data provide support for disease mechanisms wherein GPT2 plays a critical role in alanine synthesis, as well as in neuronal anaplerosis. Anaplerosis (filling - up) is the metabolic process whereby

Lonnie Shea, Ph.D. - 2020 Awardee

Biomedical Engineering
University of Michigan

"A Cell Capture Implant to Predict Acute Cardiac Allograft Rejection"

Over 36,000 solid organ transplants are conducted annually in the US, costing $30 billion. Immunosuppressive drugs protect these donor grafts from acute rejection but increase the risk of opportunistic infections and cancer, especially in pediatric transpl ant recipients, who require immune suppression for decades. As there is no method for determining which grafts will be rejected, immunosuppression is aggressively applied in a one - size- fits - all approach.

We have developed a novel implantable scaffold device to monitor recisi8 ( )06.9 (o)-(64E6BgTc -0.01-14.9 .6 (e)]TJ -0

Stephen Strittmatter, M.D., Ph.D. - 2020 Awardee

Neurology
Yale School of Medicine

“Neural Repair for Spinal Cord Injury by Axon Regeneration”

Ross Summer, M.D. - 2020 Awardee

Medicine
Thomas Jefferson University

“Preventing De - condensation of Nascent Chromatin in the Treatment of Pulmonary Fibrosis”

Allan Brasier, M.D. - 2019 Awardee

Medicine
University of Wisconsin - Madison

“Targeting the Myofibroblast Epigenome for First - In- Class Treatmenit8 w (it8 dh)1 BT /pTd [(C)-hd [(U)-r7 (922.2(it8)-26

John Bushweller, Ph.D. - 2019 Awardee

Molecular Physiology and Biological Physics
University of Virginia

“RUNX1-

Lin Guo, Ph.D. - 2019 Awardee

Biochemistry and Molecular Biology
Thomas Jefferson University

“Developing Therapeutic Agents to Rescue Neurotoxicity of FUS Aberrant Phase Transition”

It is now universally appreciated that accumulation of misfolded proteins, which can acquire alternative proteotoxic states, causes a series of deleterious molecular events resulting in numerous lethal neurodegenerative diseases. Among these, amyotrophic l ateral sclerosis

Yogendra Kanthi, M.D. - 2019 Awardee

Internal Medicine, Division of Cardiovascular Medicine
University of Michigan

"Combating Venous Thrombo - Inflammation with Precision Bio - functional Therapies"

Venous thromboembolism (VTE), comprised of deep venous thrombosis and secondary pulmonary embolism, affects 900,000 people and is the third leading cause of cardiovascular death. Current therapies are limited to anticoagulation, which carries significant b leeding risks and don't address the inflammatory processes that initiate and propagate VTE. Therapies that directly target inflammatory processes would address the underlying catalyst of disease and have the potential to fundamentally transform the treatment of VTE.

A distinguishing feature of VTE is the recruitment of leukocytes to the thrombus. We discovered that this is de(h)1.7 (1T5 v)1/14 (d)8.8 (n24.3 (o)-20.2(91)1.7 (11)-14.2a52 (c)7.2 (o))-17.90.7 (eu)240(f)-1 6.

Chaitan Khosla, Ph.D. - 2019 Awardee

Chemistry
Stanford University

"A Novel Therapeutic Approach for Major Depressive Disorder"

Major depressive disorder (MDD) is among the leading causes of illness and disability worldwide. It is a severe and life-threatening disease. Childhood trauma and insulin resistance are known risk factors for MDD. The pathophysiology of MDD remains poorly understood, and there is a serious dearth of new druggable biological targets to guide the development of improved therapeutics. This proposal outlines a research plan to identify a fundamentally new drug therapy for MDD.

In rodent models of depression, the biomolecule acetyl-carnitine promotes a rapid anti-depressant response. Endogenous acetyl-carnitine levels are also lower in the plasma of patients with MDD compared with age- and sex-matched healthy controls. Several older studies involving human subjects have also reported neurophysiological and neuropharmacological effects of oral acetyl-carnitine, although these studies were either uncontrolled or under-powered. While not a typical drug-like substance, acetyl-carnitine has properties that make it a promising lead for our purposes. It is classified as a Generally Regarded as Safe (GRAS) substance by the FDA, and is orally bioavailable, while also being transported across the blood-brain barrier.

Our goals for this Catalyst Research Award are to assemble a multidisciplinary team of researchers with relevant backgrounds in chemistry, neuropharmacology, and psychiatry to launch a comprehensive drug discovery effort aimed at optimizing the activity of acetyl-carnitine in MDD. Our technical aims are to: (1) Implement robust in vitro and in vivo assays based on our model for the mode of action of acetyl-carnitine in MDD; (2) Design and synthesize acetyl-carnitine analogs, and test them in the above assays; and (3) Compare the activity of the most promising analog vesus acetyl-carnitine in a rodent model of depression. If successful, these team-building and technical achievements will pave the way for a more intensive pursuit of a safe and effective drug candidate for MDD therapy under the Transformational Research Award Program.

Jeannette Messer, D.V.M., Ph.D. - 2019 Awardee

Inflammation and Immunity
The Cleveland Clinic

"Crafting New Weapons for the Fight Against Infectious Diseases"

Infectious diseases caused by bacteria, viruses, fungi, and single- cell parasites kill millions of people worldwide, each year. Although these microorganisms are very different in many ways, the first step of infection with each of them is microbial adher ence to human or host tissues. Regardless of the microorganism, adherence essentially always involves binding between a protein on the surface of the microbe and a carbohydrate ligand on the host. Without this binding, microbes cannot infect host tissues and cause disease. It is clear that

Jon Parquette, Ph.D. - 2019 Awardee

Chemistry and Biochemistry
The Ohio State University

"Antioxidant Nanoscaffold Technology for Combinatorial Treatment of Diabetes"

Diabetic patients often develop neuropathies and have a greater risk for vascular or Alzheimer's dementia compared to those without this disease. The epidemic of these diseases indicates a need to improve the treatment of diabetes and reduce neurodegenerat ion. Patients with diabetes must use increasingly higher doses of therapeutics over time, which increases the risk for side effects. These side- effects are partly a consequence of the fact that insulin is a growth factor that recruits glucose for anabolic processes. Consequently, diabetic patients treated with insulin are prone to weight gain and dyslipidemia that increases the risk of cardiovascular mortality. In this work, a series of short peptides are being designed and synthesized, which undergo self - assembly into nanofibers upon exposure to physiological conditions. The positive charge on the nanofibers electrostatically binds negatively charged molecules, including insulin, and can concomitantly bind to critical receptors, i.e. LepR, to stimulate gluc ose uptake as a cytokine.

Anne Rowley, M.D. - 2019 Awardee

Pediatrics
Ann & Robert H. Lurie Children's Hospital of Chicago

“A New Human Hepacivirus as an Etiologic Agent of Kawasaki Disease”

Kawasaki Disease (KD) is the leading cause of acquired heart disease in children in developed nations. KD can result in coronary artery aneurysms that can lead to lifelong heart disease, myocardial infarction, and death. The clinical and epidemiologic feat ures support an infectious etiology in genetically susceptible children, but the cause has eluded more than 50 years of study. Delayed and missed diagnoses increase the risk of coronary artery aneurysms. The development of urgently needed diagnostic tests and improved therapies are dependent upon identifying the etiology. In this study, we propose to determine if a new human hepacivirus is an etiologic agent of KD. Recent studies have shown that 1 - 2 weeks after infection with any specific pathogen, 75% of p eripheral blood plasmablasts target that specific infection. In preliminary studies, we analyzed the peripheral blood plasmablast response at 1- 3 weeks after KD fever onset using single cell RT - PCR and made 61 monoclonal antibodies (Mab) from these plasmab lasts. We used these Mab to determine their target antigens. We found that 33/61 Mab, derived from 9/11 KD patients, identify intracytoplasmic virus - like inclusion bodies (ICI) in ciliated bronchial epithelium of KD children but not infant controls. Using a viral peptide discovery array and/or ELISA, we found that 6 of the 33 (18%) Mab, derived from 3 KD patients with coronary artery aneurysms, recognize multiple similar peptides of hepacivirus non -

Sarah Slavoff, Ph.D. - 2019 Awardee

Chemistry
Yale School of Medicine

"Targeted Degradation of a Melanoma Transcription Factor"

Melanoma, the deadliest skin cancer, was diagnosed in approximately 90,000 Americans and led to over 9,000 deaths in 2019. While BRAF inhibitors and checkpoint inhibitors have revolutionized treatment of metastatic melanoma, these therapies are limited by rapid development of resistance and low response rates, respectively. New therapies, and entirely novel drug targets, are therefore critically needed, ideally targeting not only melanomas but also congenital giant nevi that are precursors to melanomas and are currently removed by repeated surgery. An innovative approach originally developed by Prof. Craig Crews at Yale University, leverages molecules called proteolysis targeting chimeras, or PROTACs, to degrade – rather than inhibit – previously "undruggabl e" classes of proteins, including transcription factors. PROTACs also have the potential to enhance immunotherapy because the proteolytic peptides arising from target protein degradation are immunogenic. In this work, we propose development of PROTACs targ eting a melanocyte - specific transcription factor, SOX10, that is required for melanoma cell, normal melanocyte and nevus cell proliferation. PROTACs consist of a ligand to the protein of interest, a flexible linker, and an E3 ubiquitin ligase ligand. We will therefore first develop a ligand to SOX10 using two high - throughput screening approaches for which preliminary feasibility has been established (Specific Aim 1). While ligand development is in progress, we will utilize a previously reported biotechnolog y- based strategy to demonstrate that PROTAC - mediated degradation of an engineered SOX10 fusion protein kills melanoma cells (Aim 2). Finally, we will create a small PROTAC library based on our novel SOX10 ligands, from which optimal members will be selected based on their ability to induce degradation of the transcription factor, inhibit melanoma cell, normal melanocytes and nevus cell proliferation, and eliminate melanocytes from human skin tissue in organotypic culture (Aim 3). In the future, we envision that the best PROTACs identified in this study will be further optimized and elaborated into drug candidates for testing in animal models prior to translation to the clinic. More broadly, this work will advance current cutting - edge efforts to drug transcri ption factors that have been refractory to traditional inhibitor development.

Takanori Takebe, M.D., Ph.D. - 2019 Awardee

Gastroenterology, Hepatology and Nutrition
Cincinnati Children's Hospital Medical Center

“Novel Human Organoid Transplant Therapy Against Pediatric Liver Disease”

Disorders affecting urea metabolism are among the most common inborn errors of metabolism in the liver. Defects in metabolism of waste nitrogen from the breakdown of protein and other nitrogen- containing molecules lead to elevated blood ammonium level that causes neurotoxicity and can be fatal. Currently, low protein diets and liver transplantation are the only available therapies for disord2143 (o)-20.2 (nd)7.4 .4 (r)1 (d)c (i)-7s(s4 (r)-9d)7./17.8 (.1 (n)-5

Hongwei Yao, Ph.D. - 2019 Awardee

Christopher Hadad, Ph.D. - 2018 Awardee

Chemistry and Biochemistry
The Ohio State University

"Development of a Novel Treatment for Organophosphorus Chemical Nerve Agent Poisoning"

Many organophosphorus (OP) compounds are chemical warfare agents or pesticides and are significant threats to human society as over 200,000 people die every year from OP exposure. These chemical agents inhibit the enzyme acetylcholinesterase (AChE) in the human body, and upon exposure to OP pesticides, AChE undergoes an aging process, rendering current oxime treatment to be ineffective. Our team has recently demonstrated the first small drug-like molecules that can reverse the effects of aging for AChE in vitro, thereby fully reversing the effects of OP exposure. The objective for this proposal is to expand on these alkylating compounds to improve selectivity, binding and reactivity with the

Oliver Jonas, Ph.D. - 2018 Awardee

Anthony Koleske, Ph.D. - 2018 Awardee

Molecular Biophysics and Biochemistry
Yale School of Medicine

"Modulators of TRIO Guanine Nucleotide Exchange Factor Activities as New Therapeutics for Schizophrenia, Autism, and Related Disorders"

Schizophrenia, autism spectrum disorder, bipolar disease, and epilepsy are genetically related and highly debilitating neurodevelopmental disorders that affect >3% of the world population. Current therapies for these diseases do not ameliorate all symptoms and have not improved in efficacy in decades. Our goal is to develop better drugs to treat these disorders.

TRIO is among the genes most frequently impacted by heterozygous rare damaging genetic variants in individuals with schizophrenia, autism, and related disorders. TRIO contains two guanine nucleotide exchange factor (GEF1 and GEF2) domains which catalyze GTP exchange on the Rac1 and RhoA GTPases, respectively, to regulate synapse development and function. Most of the disorder-associated mutations in TRIO are heterozygous and disrupt the GEF1 and/or GEF2 activities. Deletion of one TRIO allele in mice compromises normal brain development and function. Together, these data strongly implicate reduced TRIO function as a causative factor in neurodevelopmental disorders.

Nicholas Leeper, M.D. - 2018 Awardee

Surgery
Stanford University

"Precision Nanotherapies for Cardiovascular Disease"

Atherosclerosis is the process underlying heart attack and stroke. Despite recent advances, atherosclerotic cardiovascular disease (CVD) remains the leading cause of death in the United States. Most current therapies are directed against cardiovascular ris k factors (such as hypertension and elevated cholesterol levels). However, much of the population's risk of developing disease occurs independently of traditional risk factors. Therapies that directly target the plaque would instead address the root cause of disease and have the potential to fundamentally transform how CVD is treated.

A characteristic feature of the atherosclerotic plaque is the pathological accumulation of diseased and dying cells in the necrotic core. We discovered that this phenomenon is driven by the marked upregulation of a key 'don't eat me' molecule known as CD47 . This renders vascular cells 'inedible' and resistant to 'efferocytosis' (programmed cell removal). We showed that systemic delivery of anti - CD47 antibodies (Ab) could reactivate efferocytosis within the lesion, thus dramatically reducing plaque developme nt and vascular inflammation. However, systemic antibody - based therapy also caused off - target clearance of red blood cells. This induces an anemia which represents a critical roadblock in the tran10.9 (s)-e0.5 (r)-14.95cna 7nscean dnHo r

Aaron Ring, M.D., Ph.D. - 2018 Awardee

Immunobiology
Yale University

“Breaking the Vicious Cycle of Necroptosis and Inflammation with a Therapeutic “Molecular Shroud””

Sepsis is a syndrome characterized by a maladaptive immune response to infection that leads to hypotension, organ dysfunction, and death. It is a common disease with an exceedingly high fatality rate and was deemed a global health priority by the World Hea lth Organization in 2017. Despite decades of clinical and basic science research, there are no effective treatment options for sepsis other than early er ayn(o)19.6 (t)3 (h)0.5 (t)rM (h)14.2 (n)10.8m..SF36.4 11.8 (z0.

Anita Shukla, Ph.D. - 2018 Awardee

School of Engineering
Brown University

“Combating the Global Threat of Antibiotic Resistance with Bacteria - Triggered Biomaterials”

Antibiotic resistance is a growing health threat worldwide. Although microbes evolve resistance mechanisms inherently, resistance can be exacerbated by prophylactic and systemic antibiotic administration and lengthy exposures to antibiotics at sub - inhibito ry concentrations. Localized drug delivery has the potential to provide rapid antibacterial therapy at the site of an infection, while preventing offsite toxicity and reducing susceptibility to resistance. However, most currently existing local antibiotic delivery devices

John Tilton, M.D. - 2018 Awardee

Department of Nutrition
Case Western Reserve University

"Treatments and Cures for Genetic Diseases Caused by Premature Termination Codons"

Premature termination codons (PTCs) are genetic mutations that result in truncated and inactive proteins. The World Health Organization estimates that over 10,000 human diseases are caused by mutations within a single gene, and 10 - 25% of these are believed to be the caused by PTCs. While individually rare, collectively genetic diseases caused by PTCs affect many patients and their families and represent a significant burden to the health care system. Many of these diseases result in severe developmental or neurological defects or are lethal in infancy and early childhood, and almost none have effective treatments or cures. Several treatment and curative modalities for PTCs have been proposed. Suppressor tRNAs are engineered tRNAs that recognize the terminati on codon but add the appropriate amino acid instead of terminating the chain, enabling read - through of the PTC and the producing the functional protein. CRISPR - Cas9 guided single base editors can change individual base pairs within DNA, correcting the unde rlying mutations and potentially curing these diseases. Both approaches have been effective in vitro but are limited by the lack of an in vivo - compatible platform to deliver these therapies to tissues within patients. The objective of this study is to dete rmine whether our novel, rationally engineered nanoscale PrOtein Delivery (nanoPOD) platform can deliver suppressor tRNAs (Aim 1) or CRISPR - Cas9 guided single base editors (Aim 2) in vivo and enable read - through or correction of PTCs. We will employ a powe rful mouse model that contains a DNA sequence – including a PTC – from patients suffering from glycogen storage disease type 1a (GSD1a) upstream of red - shifted lucre ttoboaed

Wei Xu, Ph.D. - 2018 Awardee

Oncology
University of Wisconsin - Madison

"Development of Novel Agents for Treatment of Endocrine Resistant Breast Cancer"

Approximately 70% of breast cancers express estrogen receptor alpha (ER α) and are, therefore, treated with endocrine therapies. Although many patients benefit from Tamoxifen or aromatase inhibitors (AIs) in adjuvant and metastatic settings, approximately 50% of responsive tumors eventually relapse due to the development of res istance. One emerging mechanism of resistance is the clonal evolution of mutations in a "hotspot" within the ligand - binding domain (LBD) of ESR1, the gene encoding ER α. ESR1 mutations occur in 10- 20% of patients with metastatic ER α- positive disease who received endocrine therapies. The mutations led to ligand - independent ER α activity that promotes tumor growth and metastasis, and reduced efficacy of ER α antagonists resulting in endocrine resistance. The mutant ER α proteins ar e also more resistant to selective estrogen receptor degraders (SERDs) such as faslodex. AZD9496 and GDC - 0810, two newly developed, orally bioavailable selective estrogen receptor degraders (SERDs) are in clinical trials. However, both displayed mild estro gen activity in endometrial cells and increased uteri weight in rat models, raising the concern that the mixed agonist/antagonist activity may increase risk of endometrial cancer. Thus, our goal is to identify a new class of ERα blocker to treat metastatic, ER α- expressing breast cancer.

This study builds on our recent discovery of a natural plant product, Diptoindonesin G (Dip G) that significantly decreases ER α protein levels and, importantly, is insensitive to ESR1 mutations. We showed that Dip G acts via a mechanism distinct from all known endocrine - therapy agents. Instead of binding to ER α, the direct target of Dip G is CHIP/STUB1, an E3 ubiquitin ligase that controls ER α stability. Consequently, Dip G can degrade mutant ER α more effectively than faslodex within the therapeutic window. We hypothesize that Dip G and its analogues will be effective for treating mutant ER α expressing, endocrine- resistant tumors. This application will focus on the synthesis of novel Dip G analogues to determine the basic pharmacophore of Dip G and improve the potency and pharmacological properties of Dip G (Aim 1), which will be performed by co -